



A GUIDE TO

Post-Operative Care Tibial Tuberosity Advancement (TTA) Surgery



Post-operative care and your responsibility as the owner

Your dog has just had major orthopaedic surgery and strict adherence to the following post-operative instructions will ensure the best possible chance of an uncomplicated recovery.

NSAID pain k	iller:		
Please give:			
	Tablet(s)	Times a day	Number of days

Other:			
Please give:			
	Tablet(s)	Times a day	Number of days

Medication

Please administer all medicines given to you at discharge. Occasionally dogs will develop vomiting or diarrhoea whilst taking medication (pain relief or antibiotics). Please contact us if this happens.

Opioid:			
Please give:			
	Tablet(s)	Times a day	Number of days

Antibiotic:			
Please give:			
	Tablet(s)	Times a day	Number of days

Wounds



A pad may be covering the wound at the time of discharge from the hospital. This can be removed after two days or before if soiled. Usually there will be no external stitches as the stitches are buried under the skin and will dissolve. There should be no discharge coming from the wound. Please inform us if the wound starts to discharge or if it looks excessively swollen. Please do not apply anything to the wound as it can cause irritation.

We normally discharge patients with a buster collar. This is to prevent your dog from licking the wound and making it sore. please leave the buster collar on for 2 weeks. If an infection were to get in through the wound, this could result in further surgery.

We recommend you apply ice packs for 10 to 15 minutes several times daily during the first few days following surgery, to reduce swelling and improve comfort. Regular, gentle massage (sweeping motions from ankle to hip) may help dissipate any oedema fluid. Massage techniques will be advised at discharge. Your vet will talk you through this if this is something your dog requires.

Re-examination appointments



You should see your own vet 5 days after surgery to ensure all is progressing as planned.

If there are stitches present these will need removing at your own yet at 10-14 days

We will ask to see you and your pet at 6-8 weeks post-surgery. This recheck will involve x-rays to assess the implants and the healing of the bones. You will need to fast your pet in the morning before this appointment. If everything is progressing well, exercise can be increased further, which the surgeon will discuss with you at discharge.

If you have any queries, we are always happy to answer them

PARAGON VETERINARY REFERRALS

Paragon Veterinary Referrals, Paragon Business Village, Red Hall Crescent, Wakefield WF1 2DF

T 01924 908333 W paragonreferrals.co.uk

f #paragonveterinaryreferrals

Committed to excellence

Post-operative care



- It is better to be safe than sorry so if you are not sure about anything please contact us.
- If you are provided with pain killers please follow the instructions as directed and inform us if there is any vomiting
- Your dog should be kept confined (ideally to a large cage or alternatively a single room with non-slip flooring) to restrict activity. Confinement should be maintained at all times for the first six weeks following surgery.

We advise the following exercise programme after the operation:



• Week 1 and 2 after the operation:

We allow a maximum of two 5 minute lead walks to allow toileting only. When in the garden you should keep your dog on a short lead to prevent explosive activity.

- Week 3 and 4 after the operation:
 Two 10 minute lead walks on a short lead.
- Week 5 to 6 after the operation:
 Two 15 minute lead walks only.
- You may notice stiffness after rest as the exercise progresses. As long as this is short lived there is nothing to worry about. However, you must contact us if you notice any persistent lameness or a sudden deterioration of your dog's lameness.
- Hydrotherapy is beneficial (but not essential)
 to recovery but should only be performed in a
 centre with qualified personnel. Hydrotherapy
 may commence at three to four weeks postsurgery in a water treadmill, or at four to six
 weeks following surgery in a swimming pool.
 Any activities prior to confirmed healing of bone
 will introduce risk e.g. car travel to and from
 hydrotherapy, slipping/falling.

Signed	
Owner:	
Veterinary Surgeon:	
Date:	